

CHEM 1515.001 – 1515.006  
Exam I  
John I. Gelder  
September 12, 2001




Name \_\_\_\_\_  
TA's Name \_\_\_\_\_  
Lab Section \_\_\_\_\_

### INSTRUCTIONS:

1. This examination consists of a total of 8 different pages. The last three pages include a periodic table, a table of vapor pressures for water, a solubility table and some useful equations. All work should be done in this booklet.
2. PRINT your name, TA's name and your lab section number now in the space at the top of this sheet. DO NOT SEPARATE THESE PAGES.
3. Answer all questions that you can and whenever called for show your work clearly. Your method of solving problems should pattern the approach used in lecture. You do not have to show your work for the multiple choice or short answer questions.
4. No credit will be awarded if your work is not shown in problems 4, 5 and 7.
5. Point values are shown next to the problem number.
6. Budget your time for each of the questions. Some problems may have a low point value yet be very challenging. If you do not recognize the solution to a question quickly, skip it, and return to the question after completing the easier problems.
7. Look through the exam before beginning; plan your work; then begin.
8. ~~Relax~~ and do well.

	Page 2	Page 3	Page 4	Page 5	TOTAL
SCORES	<u>        </u> (26)	<u>        </u> (27)	<u>        </u> (26)	<u>        </u> (21)	<u>        </u> (100)

(12) 1. Write the chemical formula(s) of the product(s) and balance the following reactions. Identify all products phases as either (g)as, (l)iquid, (s)olid or (aq)ueous. Soluble ionic compounds should be written in the form of their component ions.




(4) 2a. Write the ionic and net ionic chemical equations for 1a) or 1b).

Ionic equations

Net Ionic equation

(10) 3. Identify the intermolecular attractive forces(s) present in the following substances. If more than one intermolecular force, indicate which is the most important.



- (17) 4a. The equilibrium vapor pressure for carbon disulfide,  $\text{CS}_2$ , at  $10^\circ\text{C}$  is 192 mmHg. Predict the phase(s) present at  $10^\circ\text{C}$  if 14.0 g of  $\text{CS}_2$  are placed into a 10.00 L container (previously evacuated). Support your answer with an explanation, and any important calculations.
- b. What phase(s) are present if the volume of the container were twice as large at the same temperature? (Assume no loss of  $\text{CS}_2$ )
- (10)5a. The vapor pressure of cyclohexane at  $20^\circ\text{C}$  is 78 mmHg and at  $60^\circ\text{C}$  it is 390 mmHg. Calculate the heat of vaporization,  $\Delta H^\circ_{\text{vap}}$  for cyclohexane.

- (16) 6. The boiling point of the first two binary hydrogen compounds in Group IV and V are shown in the Table below:

Compound	Boiling Point (°C)
CH <sub>4</sub>	-164
SiH <sub>4</sub>	-112
H <sub>2</sub> O	100
H <sub>2</sub> S	-61

Explain why CH<sub>4</sub> has a lower boiling point compared to SiH<sub>4</sub>, but H<sub>2</sub>O has a higher boiling point compared to H<sub>2</sub>S?

- (10) 7. Tungsten crystallizes in a face-centered cubic unit cell. The atomic radius of a tungsten atom is 0.137 nm. Calculate the density, in g cm<sup>-3</sup>, of the unit cell.

(8) 8. For each of the following solids identify the type of attractive force(s) that exists?

a)  $\text{SiO}_2$

b)  $\text{N}_2\text{O}_4$

c)  $\text{Mo}$

d)  $\text{MgF}_2$

(13) 9a. Name the two phases involved in evaporation.

b) Write a chemical equation to symbolically represent evaporation for a substance of your choosing.

c) Is evaporation endothermic or exothermic?

d. From a particulate level (atomic level) how does evaporation occur?

## Periodic Table of the Elements

	IA															VIIIA		
1	1 <b>H</b> 1.008															2 <b>He</b> 4.00		
2	3 <b>Li</b> 6.94	4 <b>Be</b> 9.01											5 <b>B</b> 10.81	6 <b>C</b> 12.01	7 <b>N</b> 14.01	8 <b>O</b> 16.00	9 <b>F</b> 19.00	10 <b>Ne</b> 20.18
3	11 <b>Na</b> 22.99	12 <b>Mg</b> 24.30	IIIB	IVB	VB	VIB	VIIB	VIII		IB	IIB	13 <b>Al</b> 26.98	14 <b>Si</b> 28.09	15 <b>P</b> 30.97	16 <b>S</b> 32.06	17 <b>Cl</b> 35.45	18 <b>Ar</b> 39.95	
4	19 <b>K</b> 39.10	20 <b>Ca</b> 40.08	21 <b>Sc</b> 44.96	22 <b>Ti</b> 47.88	23 <b>V</b> 50.94	24 <b>Cr</b> 52.00	25 <b>Mn</b> 54.94	26 <b>Fe</b> 55.85	27 <b>Co</b> 58.93	28 <b>Ni</b> 58.69	29 <b>Cu</b> 63.55	30 <b>Zn</b> 65.38	31 <b>Ga</b> 69.72	32 <b>Ge</b> 72.59	33 <b>As</b> 74.92	34 <b>Se</b> 78.96	35 <b>Br</b> 79.90	36 <b>Kr</b> 83.80
5	37 <b>Rb</b> 85.47	38 <b>Sr</b> 87.62	39 <b>Y</b> 88.91	40 <b>Zr</b> 91.22	41 <b>Nb</b> 92.91	42 <b>Mo</b> 95.94	43 <b>Tc</b> (98)	44 <b>Ru</b> 101.1	45 <b>Rh</b> 102.9	46 <b>Pd</b> 106.4	47 <b>Ag</b> 107.9	48 <b>Cd</b> 112.4	49 <b>In</b> 114.8	50 <b>Sn</b> 118.7	51 <b>Sb</b> 121.8	52 <b>Te</b> 127.6	53 <b>I</b> 126.9	54 <b>Xe</b> 131.3
6	55 <b>Cs</b> 132.9	56 <b>Ba</b> 137.3	57 <b>La</b> 138.9	72 <b>Hf</b> 178.5	73 <b>Ta</b> 180.9	74 <b>W</b> 183.8	75 <b>Re</b> 186.2	76 <b>Os</b> 190.2	77 <b>Ir</b> 192.2	78 <b>Pt</b> 195.1	79 <b>Au</b> 197.0	80 <b>Hg</b> 200.6	81 <b>Tl</b> 204.4	82 <b>Pb</b> 207.2	83 <b>Bi</b> 209.0	84 <b>Po</b> (209)	85 <b>At</b> (210)	86 <b>Rn</b> (222)
7	87 <b>Fr</b> (223)	88 <b>Ra</b> 226.0	89 <b>Ac</b> 227.0	104 (261)	105 (262)	106 (263)												

Lanthanides	58 <b>Ce</b> 140.1	59 <b>Pr</b> 140.9	60 <b>Nd</b> 144.2	61 <b>Pm</b> (145)	62 <b>Sm</b> 150.4	63 <b>Eu</b> 152.0	64 <b>Gd</b> 157.2	65 <b>Tb</b> 158.9	66 <b>Dy</b> 162.5	67 <b>Ho</b> 164.9	68 <b>Er</b> 167.3	69 <b>Tm</b> 168.9	70 <b>Yb</b> 173.0	71 <b>Lu</b> 175.0
Actinides	90 <b>Th</b> 232.0	91 <b>Pa</b> 231.0	92 <b>U</b> 238.0	93 <b>Np</b> 237.0	94 <b>Pu</b> (244)	95 <b>Am</b> (243)	96 <b>Cm</b> (247)	97 <b>Bk</b> (247)	98 <b>Cf</b> (251)	99 <b>Es</b> (252)	100 <b>Fm</b> (257)	101 <b>Md</b> (258)	102 <b>No</b> (259)	103 <b>Lr</b> (260)

## Useful Information

$$PV = nRT$$

$$\ln\left(\frac{vp_2}{vp_1}\right) = -\frac{\Delta H^\circ_{\text{vap}}}{R}\left(\frac{1}{T_2} - \frac{1}{T_1}\right)$$

$$1 \text{ nm} = 10^{-9} \text{ m}$$

$$\text{edge length (l)} = 2r$$

$$6.023 \times 10^{23}$$

$$R = 0.0821 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} = 8.314 \frac{\text{J}}{\text{mol}\cdot\text{K}}$$

$$\text{density of H}_2\text{O} = 1.00 \frac{\text{g}}{\text{cm}^3}$$

$$1 \text{ pm} = 10^{-12} \text{ m} \quad 1 \text{ atm} = 760 \text{ mmHg}$$


$$\text{edge length (l)} = 2\sqrt{2} \cdot r \quad \text{edge length (l)} = \frac{4r}{\sqrt{3}}$$

Temperature (°C)	Vapor Pressure(mmHg)	Temperature (°C)	Vapor Pressure(mmHg)
-5	3.2	50	92.5
0	4.6	55	118.0
5	6.52	60	149.4
10	9.20	65	187.5
15	12.8	70	233.7
20	17.5	75	289.1
25	23.8	80	355.1
30	31.8	85	433.6
35	42.1	90	525.8
40	55.3	95	633.9
45	71.9	100	760

### Solubility Table

<u>Ion</u>	<u>Solubility</u>	<u>Exceptions</u>
NO <sub>3</sub> <sup>-</sup>	soluble	none
ClO <sub>4</sub> <sup>-</sup>	soluble	none
Cl <sup>-</sup>	soluble	except Ag <sup>+</sup> , Hg <sub>2</sub> <sup>2+</sup> , *Pb <sup>2+</sup>
I <sup>-</sup>	soluble	except Ag <sup>+</sup> , Hg <sub>2</sub> <sup>2+</sup> , Pb <sup>2+</sup>
SO <sub>4</sub> <sup>2-</sup>	soluble	except Ca <sup>2+</sup> , Ba <sup>2+</sup> , Sr <sup>2+</sup> , Hg <sup>2+</sup> , Pb <sup>2+</sup> , Ag <sup>+</sup>
CO <sub>3</sub> <sup>2-</sup>	insoluble	except Group IA and NH <sub>4</sub> <sup>+</sup>
PO <sub>4</sub> <sup>3-</sup>	insoluble	except Group IA and NH <sub>4</sub> <sup>+</sup>
-OH	insoluble	except Group IA, *Ca <sup>2+</sup> , Ba <sup>2+</sup> , Sr <sup>2+</sup>
S <sup>2-</sup>	insoluble	except Group IA, IIA and NH <sub>4</sub> <sup>+</sup>
Na <sup>+</sup>	soluble	none
NH <sub>4</sub> <sup>+</sup>	soluble	none
K <sup>+</sup>	soluble	none

\*slightly soluble



- (9) 8a. Describe what happened in the “ammonia fountain” experiment which was demonstrated in class. Recall a clamped 5.0 L round-bottom flask containing  $\text{NH}_3(g)$  was suspended above a second 5.0 L round-bottom flask filled with  $\text{H}_2\text{O}(l)$  to which had been added a few drops of phenolphthalein. The two flasks were connected by a glass tube which protruded through the rubber stoppered flask containing the ammonia. The flask containing water was open to the atmosphere. A small pipet bulb, containing water, was connected via a second glass rod which protruded through the rubber stoppered flask containing the ammonia. To begin the experiment a small amount of water was injected into the flask containing the  $\text{NH}_3(g)$  by a slight squeeze of the pipet bulb.



- b. Briefly provide an explanation for the observed behavior.
- c. Do you think the solution process was exothermic or endothermic?
- d. Using Lewis structures, draw a picture (and label) depicting the intermolecular interactions in the solution at the completion of the demonstration (after the fountain stopped.)