

CHEM 1515.001
Exam II
John II. Gelder
March 8, 2001



Name _____
TA's Name _____
Lab Section _____

INSTRUCTIONS:

1. This examination consists of a total of 8 different pages. The last three pages include a periodic table, a table of vapor pressures for water, a solubility table, and a table of ionic structures and packing. All work should be done in this booklet.
2. PRINT your name, TA's name and your lab section number now in the space at the top of this sheet. DO NOT SEPARATE THESE PAGES.
3. Answer all questions that you can and whenever called for show your work clearly. Your method of solving problems should pattern the approach used in lecture. You do not have to show your work for the multiple choice or short answer questions.
4. No credit will be awarded if your work is not shown in problems 4, 5 and 9.
5. Point values are shown next to the problem number.
6. Budget your time for each of the questions. Some problems may have a low point value yet be very challenging. If you do not recognize the solution to a question quickly, skip it, and return to the question after completing the easier problems.
7. Look through the exam before beginning; plan your work; then begin.
8. **Relax** and do well.

	Page 2	Page 3	Page 4	Page 5	TOTAL
SCORES	<u> </u> (25)	<u> </u> (26)	<u> </u> (20)	<u> </u> (29)	<u> </u> (100)

(9) 1. Write the chemical formula(s) of the product(s) and balance the following reactions. Identify all products phases as either (g)as, (l)iquid, (s)olid or (aq)ueous. Soluble ionic compounds should be written in the form of their component ions.



(4) 2. Write the ionic and net ionic chemical equations for 1b).

1b)

Ionic equation:

Net Ionic equation:

(12) 3. When a liquid solute is added to a liquid solvent we discussed three steps important to the solution process as it related to the enthalpy of the solution.


a) List the three steps and indicate whether the step is exothermic or endothermic;

b) In terms of these three steps explain why a particular solute does not dissolve in a particular solvent.

- (10) 4. A certain metal has a density of 10.200 g mL^{-1} at $25 \text{ }^\circ\text{C}$. It crystallizes in a body-centered cubic unit cell with an atomic radius of $1.36 \times 10^{-8} \text{ cm}$. Calculate the molar mass of the element and identify the substance.

- (10) 5. In magnesium oxide the ionic radius of Mg^{2+} is 0.65 \AA and the ionic radius of O^{2-} is 1.40 \AA . Assuming the oxide ion defines the unit cell and the cation occupies holes in the lattice, describe a probable structure for the ionic compound.

- (6) 6. Give the name or draw the Lewis structure for each of the following compounds.

	3-ethyl-3-methylpentane	2,5,5-trimethyl-3-hexyne
--	-------------------------	--------------------------

(10) 7. Draw and name four of the structural isomers for C_6H_{10} .

(10) 8. Which member of the following pairs is more soluble in water? Provide a brief explanation supporting your choice. Also provide a brief explanation why you ruled out the other choice.

a) HCl or C_4H_9Cl

b) CH_3NH_2 or $(CH_3)_3N$

- (29) 9. A solution of formic acid, HCOOH , is prepared by 54.0 g of formic acid in enough water to make 250 mLs of solution.
- calculate the molarity of the solution;
 - the solution described above is also 20.0 % by weight formic acid. Calculate the mole fraction of formic acid in the solution;
 - assuming formic acid is a nonvolatile, nonelectrolyte calculate the freezing point of this solution;
 - formic acid is actually a weak acid. Suggestion a reasonable experimental freezing point (do not calculate) for this solution? Explain your answer.
 - Calculate the density of the solution.

Periodic Table of the Elements

	IA																VIII A	
1	1 H 1.008																	2 He 4.00
2	3 Li 6.94	IIA	4 Be 9.01									IIIA	IVA	VA	VIA	VIIA	10 Ne 20.18	
3	11 Na 22.99	12 Mg 24.30										5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	18 Ar 39.95	
4	19 K 39.10	20 Ca 40.08	IIIB	IVB	VB	VIB	VIIB	VIII		IB	IIB	13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	36 Kr 83.80	
5	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
6	55 Cs 132.9	56 Ba 137.3	57 La 138.9	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po (209)	85 At (210)	86 Rn (222)
7	87 Fr (223)	88 Ra 226.0	89 Ac 227.0	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)									

Lanthanides	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm (145)	62 Sm 150.4	63 Eu 152.0	64 Gd 157.2	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0
Actinides	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.0	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)

Useful Information

$$T = ikm \quad k_f(\text{H}_2\text{O}) = 1.86 \frac{^\circ\text{C}}{\text{m}} \quad k_b(\text{H}_2\text{O}) = 0.512 \frac{^\circ\text{C}}{\text{m}}$$

$$R = 0.0821 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} = 8.314 \frac{\text{J}}{\text{mol}\cdot\text{K}}$$

$$P_{\text{solution}} = \text{solvent } P^\circ_{\text{solvent}}$$

$$\text{edge length (l)} = 2r$$

$$\text{edge length (l)} = 2\sqrt{2} \cdot r$$

$$\text{edge length (l)} = \frac{4r}{\sqrt{3}}$$

$$\text{density of H}_2\text{O} = 1.00 \frac{\text{g}}{\text{cm}^3}$$

$$6.023 \times 10^{23}$$

Temperature (°C)	Vapor Pressure(mmHg)	Temperature (°C)	Vapor Pressure(mmHg)
-5	3.2	50	92.5
0	4.6	55	118.0
5	6.52	60	149.4
10	9.20	65	187.5
15	12.8	70	233.7
20	17.5	75	289.1
25	23.8	80	355.1
30	31.8	85	433.6
35	42.1	90	525.8
40	55.3	95	633.9
45	71.9	100	760

Solubility Table

<u>Ion</u>	<u>Solubility</u>	<u>Exceptions</u>
NO ₃ ⁻	soluble	none
ClO ₄ ⁻	soluble	none
Cl ⁻	soluble	except Ag ⁺ , Hg ₂ ²⁺ , *Pb ²⁺
I ⁻	soluble	except Ag ⁺ , Hg ₂ ²⁺ , Pb ²⁺
SO ₄ ²⁻	soluble	except Ca ²⁺ , Ba ²⁺ , Sr ²⁺ , Hg ²⁺ , Pb ²⁺ , Ag ⁺
CO ₃ ²⁻	insoluble	except Group IA and NH ₄ ⁺
PO ₄ ³⁻	insoluble	except Group IA and NH ₄ ⁺
-OH	insoluble	except Group IA, *Ca ²⁺ , Ba ²⁺ , Sr ²⁺
S ²⁻	insoluble	except Group IA, IIA and NH ₄ ⁺
Na ⁺	soluble	none
NH ₄ ⁺	soluble	none
K ⁺	soluble	none

*slightly soluble

Simple Ionic Structures Grouped According to Anion Packing

Structure Name	Anion Packing	Coordination Number	Sites Occupied by Cations	Examples
Rock Salt	ccp	6:6 MO	all octahedral	NaCl, LiF, KBr, CdO, FeO
Zinc Blende	ccp	4:4 MO	$\frac{1}{2}$ tetrahedral	ZnS, BeO, SiC
Antifluorite	ccp	4:8 M ₂ O	all tetrahedral	Li ₂ O, sulfides
Rutile	distorted ccp	6:3 MO ₂	$\frac{1}{2}$ octahedral	TiO ₂ , GeO ₂ , MnO ₂ , OsO ₂
Perovskite	ccp	12:6:6 ABO ₃	$\frac{1}{4}$ octahedral(B)	CaTiO ₃ , SrSnO ₃
Spinel	ccp	4:6:4 AB ₂ O ₄	$\frac{1}{8}$ tetrahedral(A) $\frac{1}{2}$ octahedral(B)	MgAl ₂ O ₄ , FeAlO ₄
Cesium Chloride	simple cubic	8:8 MO	all cubic	CsCl, CsBr, CsI
Fluorite	simple cubic	8:4 MO ₂	$\frac{1}{2}$ cubic	CaF ₂ , UO ₂ , ThO ₂

Lattice Types and Radius Ratios of Cations and Anions

Radius Ratio (Cation/Anion)	Lattice Type	Coordination Number of Cation	Coordination Number of Anion
--------------------------------	--------------	----------------------------------	---------------------------------

A. 1:1 Stoichiometry of Salt (MX)

0.225 – 0.414	Zinc Blende	4	4
0.414 – 0.732	Rock salt (NaCl)	6	6
0.732 – 1.000	Cesium chloride	8	8

B. 1:2 Stoichiometry of Salt (MX₂)

0.225 – 0.414	Beta-quartz	4	2
0.414 – 0.732	Rutile (TiO ₂)	6	3
0.732 – 1.000	Fluorite (CaF ₂)	8	4